

Exponential Yield Surge: The GPL-37 Pakistan Protocol

Professional Proof-of-Concept
Report: Achieving >200% Growth
in Lemon Tree Cultivation



>200%

Document Reference: 201706 / GROWING | Field Location:
Pakistan | Agent: GPL-37 Chain Reaction Fertilizer


Field Trial Parameters & Headline Efficacy

1




Crop: Lemon Tree
Location: Pakistan

2



Compound: GPL-37
Chain Reaction Fertilizer

3




A commercial agricultural client in Pakistan reported a “miracle experience” utilizing GPL-37. By deploying a specific sequential application method, the trial resulted in a verified yield increase exceeding 200% over baseline production.

The Sequential Application Timeline



The breakthrough yield was not achieved by a single application, but by a sequenced protocol separating root and foliar absorption by exactly 15 days.

Bimodal Uptake: Systemic Foundation vs. Targeted Catalyst



Diagnostic Matrix: Application Modalities

	Aqueous Root Pour	Direct Foliar Spray
Carrier Method	Mixed in water	Direct aerial application
Labor & Friction	Standard / Low friction	High (A difficult job)
Observed Efficacy	Good (Baseline performance)	Exceptional (Far better than pouring)

Field observations confirm that while foliar spraying introduces higher operational friction, its efficacy profile dramatically outperforms standard aqueous root pouring.


The Friction vs. Reward Equation

“Suggest we spray on the tree even if it’s a difficult job.”



The high operational friction of full-canopy spraying is entirely negated by the disproportionate ROI of the resulting yield surge. Spraying is non-negotiable for maximum yield.

The 'Chain Reaction' Synthesis Framework



The 200% yield surge is not solely the result of the compound itself, but the synergistic sequence of its application. The initial root pour primes the vascular system, while the the delayed foliar spray triggers a compounding "chain reaction" of vegetative and fruiting growth.

Standardized Protocol Recommendation

1.

Initiate with water-mixed **root** infusion.

2.

Maintain strict 15-day application interval.

3.

Execute comprehensive **foliar spray** (Mandatory, despite labor difficulty).

Based on this unique application discovery, GROWING officially recognizes this bimodal sequential method as the optimal protocol for maximizing the efficacy of the GPL-37 chain reaction agent.

Play to Win. Partner with Nature.

The future of global agriculture relies on the intersection of biological science and environmental stewardship. **Growing-Up INTL Corp** provides the definitive, zero-compromise solution for sustainable crop management.

Contact & Partnership Hub

Growing-Up International Corp.
69, Minchuan Rd, Tsao Tuen, Nantou, Taiwan 54262
Tel: +886-49-2337927
Email: growing8@growing8.com
Web: www.growing8.com

